

LA English
 AB The influence of the DA D2 antagonist (-)-eticlopride on cocaine
 - and DA D2 agonist-induced behavioral effects was investigated by means
 of two series of expts., in rats. In the first 10-day series,
 coadministration of (-)-eticlopride (10 and 50 .mu.g/kg, SC) always
 potentially inhibited cocaine (15 mg/kg, IP)-induced hypermotility
 but did not modify the penile erection (PE)-enhancement produced by the
 drug at the first injection; it actually counteracted the inhibitory
 effect of subchronic cocaine on PE. In the second series,
 (-)-eticlopride, at the same doses, antagonized PE elicited by various DA
 D2 agonists at nonstereotyping doses; when, along with PE, stereotyped
 behavior was induced, only the latter was inhibited by (-)-eticlopride,
 which even increased PE.

L12 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2002 ACS

AN 1991:598533 CAPLUS

DN 115:198533

TI Use of dopamine autoreceptor agonists in the treatment of drug dependency

IN Kutter, Eberhard; Schingnitz, Guenter

PA Boehringer Ingelheim K.-G., Germany; Boehringer Ingelheim International
 G.m.b.H.

SO Eur. Pat. Appl., 13 pp.

CODEN: EPXXDW

DT Patent

LA German

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 417637	A2	19910320	EP 1990-117147	19900906
	EP 417637	A3	19920902		
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE				
	DE 3930282	A1	19910321	DE 1989-3930282	19890911
	DD 297557	A5	19920116	DD 1990-343855	19900906
	CA 2025003	AA	19910312	CA 1990-2025003	19900910
	JP 03106825	A2	19910507	JP 1990-239790	19900910
	HU 57584	A2	19911230	HU 1990-5853	19900910
PRAI	DE 1989-3930282		19890911		

AB BHT 920 (I) and SND 919 (II) and their acid addn. salts are dopamine
 autoreceptor agonists (i.e. decrease the synthesis and release of dopamine
 from cells of the mesolimbic and nigrostriatal system) and thus are useful
 in treatment of drug dependence mediated by dopamine release. By
 diminishing the pos. reinforcement of drug consumption resulting from
 dopamine release in these brain centers and the consequent euphoric inner
 reward, I and II prevent craving for the drug. The action of I and II is
 enhanced by their activity on supersensitive postsynaptic D2-dopaminergic
 receptors in dopamin-depleted chronic drug abusers, as well as by their
 central .alpha.2-adrenergic activity. I and II themselves do not induce
 dependence. Thus, in monkeys allowed to self-administer cocaine
 , the self-administration rate decreased to 0 after i.v. injection of I
 (0.1 mg/kg, twice). Pills were prepd. contg. I 50 .mu.g, lactose 38.45,
 corn starch 10.0, gelatin 1.0, and Mg stearate 0.5 mg.

=> d his

(FILE 'HOME' ENTERED AT 09:26:48 ON 11 OCT 2002)

FILE 'REGISTRY' ENTERED AT 09:26:58 ON 11 OCT 2002

E PRAMIPEXOLE/CN

L1 1 S E3

E LAMOTRIGINE/CN

L2 1 S E3

0.1 mg/kg x 75 kg
 2(7.5) 14.5 g